Illegible handwriting in medical records

F Javier Rodríguez-Vera MD Y Marín MD A Sánchez MD C Borrachero MD E Pujol MD

J R Soc Med 2002;95:545-546

SUMMARY

In clinical records many items are handwritten and difficult to read. We examined clinical histories in a representative sample of case notes from a Spanish general hospital. Two independent observers assigned legibility scores, and a third adjudicated in case of disagreement. Defects of legibility such that the whole was unclear were present in 18 (15%) of 117 reports, and were particularly frequent in records from surgical departments.

Through poor handwriting, much information in medical records is inaccessible to auditors, to researchers, and to other clinicians involved in the patient's care. If clinicians cannot be persuaded to write legibly, the solution must be an accelerated switch to computer-based systems.

INTRODUCTION

Despite the computer revolution, much information in clinical records continues to be handwritten. The originator may understand what has been written, but difficulties can arise when other parties are involved. Only a few studies, however, have been reported on the legibility of medical documents and these largely about prescriptions^{1–7}. We therefore decided to examine the legibility of case histories written on admission of patients to our hospital.

METHODS

The hospital, located in south-west Spain, has 600 beds. We obtained a representative sample by examining, on a single day, case notes from patients whose rooms had even numbers. Certain specialties—intensive care, haematology, gynaecology, paediatrics—had their own record systems and were excluded. The 'clinical history' was taken to be any document written by a clinician that included the patient's name, age, medical condition, and reason for admission. Two medical residents, recently arrived at the hospital and not involved in the admissions or recording of case notes, evaluated the legibility of the document on a score of 1–4. This classification (Box 1) has been used by others⁵. They went through a training process in order to reach a kappa concordance coefficient

E-mail: frodriguezv@sego.es

Box 1 Legibility scoring

- 1 Illegible (most or all words impossible to identify)
- 2 Most words illegible; meaning of the whole unclear
- 3 Some words illegible, but report can be understood by a clinician
- 4 Legible (all words clear)

of 0.85. A third reviewer adjudicated in case of disagreement.

RESULTS

117 case notes were examined and 18 (15%) were scored 1 or 2—i.e. they were so illegible that the meaning was unclear. Table 1, giving results for individual specialties, indicates that surgical departments performed worse than medical departments.

DISCUSSION

A weakness of this study was that it might have been skewed by the poor handwriting of just a few clinicians who were responsible for many admissions. Also, we did not distinguish between cold admissions, in which the notes might simply consist of a shorthand reminder of the outpatient consultation, and acute admissions, where a full and comprehensible history is more important. This might partly explain why medical departments scored better in this respect than surgical departments.

If 15% of case histories are illegible, does this matter? In principle, it is a source of avoidable error—for audit, research, and clinical communication^{8–10}. The remedy lies

Internal Medicine Department, Hospital Juan Ramón Jiménez de Huelva, Huelva, Spain

Correspondence to: F Javier Rodríguez-Vera, C/Arjona No 12 Esc 2 1 $^{\circ}$ A, 41001 Sevilla, Spain

Table 1 Scores for individual departments

Section	Patients (n)	Score			
		1	2	3	4
Medical					
General internal medicine	33	0	1	3	29
Cardiology	7				7
Gastroenterology	5				5
Respiratory diseases	8			2	6
Neurology	8		1	1	6
Oncology	6				6
Subtotal	67	0 (0%)	2 (3%)	6 (9%)	59 (88%)
Surgical					
Orthopaedic surgery	17	1	0	5	11
General surgery	15	5	3	2	5
Otolaryngology	4	2	1	0	1
Vascular surgery	8	0	0	0	8
Ophthalmology	3	0	2	1	0
Urology	3	0	2	0	1
Subtotal	50	8 (16%)	8 (16%)	8 (16%)	26 (52%)
Total	117	8 (7%)	10 (9%)	14 (12%	81 (70%)

either in a more conscientious approach to record-keeping, with an eye to the needs of other readers, or an accelerated move towards computer-based systems¹¹. In our view, it is time to say goodbye to manuscript in medical notes, whether legible or not.

REFERENCES

- 1 Winslow EH, Nestor VA, Davidoff SK, Thompson PG, Borum JC. Legibility and completeness of physicians' handwritten medication orders. *Heart Lung* 1997;26:158–64
- 2 Brodell RT, Helms SE, KrishnaRao I, Bredle DL. Prescription errors. Legibility and drug name confusion. *Arch Fam Med* 1997;**6**:296–8
- 3 Russell N, Morrison R, Johnson S, Robertson WO. Reducing errors by increasing legibility. West J Med 2000;173:163

- 4 Kozak EA, Dittus RS, Smith WR, Fitzgerald JF, Langfeld CD. Deciphering the physician note. J Gen Intern Med 1994;9:52–4
- 5 Berwick DM, Winickoff DE. The truth about doctors' handwriting: a prospective study. BMJ 1996;313:1657–8
- 6 Lyons R, Payne C, McCabe M, Fielder C. Legibility of doctors' handwriting: quantitative comparative study. BMJ 1998;317:863–4
- 7 Cheeseman GA, Boon N. Reputation and the legibility of doctors' handwriting in situ. Scott Med J 2001;46:79–80
- 8 Bruner A, Kasdan ML. Handwriting errors: harmful, wasteful and preventable. J Ky Med Assoc 2001;99:189–92
- 9 Hester DO. Do you see what I see? Illegible handwriting can cause patient injuries. J Ky Med Assoc 2001;99:187
- 10 Brodell RT, Helms SE, KrishnaRao I, Bredle DL. Prescription errors. Legibility and drug name confusion. Arch Fam Med 1997;6:296–8
- 11 Bradbury A. Computerized medical records: the need for a standard. J Am Med Rec Assoc 1990;61:25–37